

Rising Trends towards Herbal Contraceptives

*Dr. Meera Agarwal

**Dr Sonalika Singh Jadaun

ABSTRACT

Oral administration of aqueous extract of *Calendula officinalis* flower to male rats at the dose level of 100mg/kg body weight for 60 days did not cause body weight loss but decreased the weight of testis, epididymis, seminal vesicle and ventral prostate in a significant manner. Sperm motility as well as sperm density were reduced highly significantly which resulted in 45% negative fertility. Serum testosterone level showed highly significant reduction. Biochemical parameters like total protein and Sialic acid in testis, epididymis, seminal vesicles and ventral prostate were decreased significantly where as testicular cholesterol concentration was elevated. All the haematological parameters were in normal range when compared with the control group animals.

Keywords- *Calendula officinalis*, Antispermatogetic, Sialic acid, Sperm motility, Sperm density

INTRODUCTION

In many developing countries, traditional medicines are widely utilized in the treatment of various ailments on an empirical basis. A variety of plants have been used for the treatment of diabetes and (Upadhyay et al 2004) and male reproduction (Das et al 2004). *Calendula officinalis* belongs to the family Compositae is commonly known as 'marigold'. Extract of dried flower from *Calendula officinalis* were examined for their ability to inhibit the human immunodeficiency virus type - 1 (HIV-1) replication (Kalvatchev et al 1997). It is also traditionally being used for gastric ulcers and menstrual discomforts (Szakiel et al 2005) skin disorders, antiseptic and anti-inflammatory disease (Cordova et al 2002). Contraceptive like properties have been reported in females by local tribes of Rajasthan they use it for birth control.

The contraceptive efficacy of this plant has not been investigated scientifically hence the present investigation has been undertaken to investigate the effect of aqueous extract of *Calendula officinalis*. In this communication a pronged approach involving haematological and biochemical parameters has been used and antispermatogetic activity of *Calendula officinalis* was discussed.

MATERIAL AND METHOD

The flowers of *Calendula officinalis* were collected from University Campus identified in Department of Botany (RUBL 20102), shade dried and powdered and (100)gm dry powder was macerated in 200ml of distilled water and stayed for 36 hours at room temperature and filtered to obtain a final crude extract in the form of powder. 24.75% yield was obtained from 100 gms of

flower. This powder was dissolved in distilled water which was administered to the male rats while control group rat received equal amount of distilled water. Adult, healthy male albino rats of wistar strain 16-18 week old were selected from the inbred colony and the animal were maintained according to the guide lines for care and use of animals for scientific research (Indian National Science Academy, 2000) through out the course of investigation.

The rats were divided in two groups having 10 rats in each group.

Group 1- Vehicle treated i.e. 0.5ml/rat/day distilled water for 60 days.

Group 2- 100mg/rat/day *Calendula officinalis* (COFAq) dissolved in 0.5ml of distilled water for 60 days.

FERTILITY TEST

The mating test of control and treated groups were performed on day 55-60 using the method of W.H.O (W.H.O Protocol 1990) the females were separated for normal delivery. On 16th day of pregnancy the implantation site (normal and absorbed foetus) were recorded.

AUTOPSY

After 24 hours of last dose rats were weight and autopsied under light ether anesthesia the blood was collected from heart in pre-heparinized tubes for hematological studies and serum was also separated from non-heparinized tubes for RIA studies. The animal were autopsied, the reproductive and vital organs (testis, epididymis, seminal vesicle, ventral prostate, liver, adrenal and kidney) were taken out and trimmed free of fat and weight separately on electronic balance.

SPERM MOTILITY AND DENSITY

At autopsy, the epididymis was exposed and spermatozoa were taken out by cutting cauda epididymis for sperm motility (Srikanth et al 1999) and sperm density (Zaneveld and Polakoski 1997).

HAEMATOLOGY

Total erythrocyte Count (Schalm et al, 1975), Total leukocyte count (Lynch et al 1969), haematocrit by microhaematocrit method (Schalm et al 1975). The Haemoglobin level was estimated by cynomethanoglobin method (Makarem et al 1974) blood sugar (Astoor and King 1954) and blood urea (Varley 1969) were estimated while serum was assessed for the estimation of testosterone by Radio Immuno assay (commercial kit).

BIOCHEMISTRY

Frozen testis, epididymis, seminal vesicle and ventral prostate were used for the estimation of protein (Lowry et al 1951), glycogen (Montgomery 1957), cholesterol (Oser 1965) and Sialic acid (Warren 1959).

STATISTICAL ANALYSIS

The mean and standard error of mean (SEM) were calculated from the data obtained by the

experiment and The treated groups were compared to the control using the student's ' t' test (Ipstein and poly 1970).

RESULTS

Body and organ weight-

Oral administrations of *Calendula officinalis* flower extract (COFAq) did not cause any change in the body weight when compared to their initial body weight. However it showed significant reduction in weight of testes, epididymis, seminal vesicles and ventral prostate ($p < 0.001$) in comparison to the control group (Table -1)

Sperm dynamics and serum testosterone -

Percentage of sperm motility, sperm density were decreased significantly ($p < 0.01$) where as fertility rat was 45% negative after the administration of *Calendula officinalis* (flower) aqueous extract. Serum testosterone level was reduced significantly when compared with control group. Number of pregnant females; number of implantation sites and number of viable fetuses were also declined in G-11 (Table-2).

Tissue Biochemistry-

Total protein and Sialic acid content of testis, epididymis, seminal vesicles and ventral prostate were decreased significantly following the administration of *Calendula officinalis* (flower) aqueous extract (COFAq). Glycogen level in testis and liver reduced slightly where as cholesterol level was increased slightly (Table-3).

Blood Profiles-

Calendula officinalis showed that total erythrocytes count, total leukocyte count, haemoglobin, haematocrit, blood sugar and blood urea were in normal range (Table-4).

DISCUSSION

Oral administration of *Calendula officinalis* flower extract (COFAq) showed reduction in the weight of testes, epididymis, seminal vesicles and ventral prostate. Reduction in weight of testis and other accessory sex organs might be due to low level of androgens (Sharma and Jacob, 2001), which was reflected in decreased serum testosterone level in treated rats.

Sperm motility and density in cauda epididymis and testis were decreased which shows alteration in maturation and production of sperm (Sarkar et al 2000).

Protein content of reproductive organs were significantly was decreased due to low level of androgens (Chinoy and Bhattacharya 1997) which was confirmed in low concentration of serum testosterone.

Decreased level of Sialic acid in testis, epididymis, seminal vesicles and ventral prostate reflected loss of androgens (Gupta et al 2001). Mode of action of *Calendula officinalis* extract was through pituitary gonadal axis, which was confirmed in decreased serum testosterone level. After the administration *Calendula officinalis* extract, increased testicular cholesterol might be due to

arrest of steroidogenesis of testosterone (Gupta et al 2002) so to accumulate in the testis.

From the present study it is concluded that the oral administration of crude ethanolic extract of *Calendula officinalis* may lead to fertility control in male rats due to interfere in the testicular androgens level which arrest the process of spermatogenesis in testis without disturbing general metabolism.

***Retd. Associate Professor**

Department of Zoology, University of Rajasthan

****Assistant Professor**

Department of Zoology, S.S. Jain Subodh Girls College

BIBLIOGRAPHY

Astoor A. and King E.J., (1954): Simplified Calorimetric Blood Sugar Method Journal of Biological Chemistry 56 XIV.

Chinoy M.R.N. and Bhattacharya S., (1997): Effect of chronic administration of aluminium chloride on reproductive function of testes and accessory sex organs of male mice. Journal of Environmental Toxicology 7, 12-15.

Cordova C.A., Netlo C.A., Yunes R.A., (2002): Calendula used for the treatment of skin disorders antiseptic and anti-inflammatory disease. Redox Reproduction 7, 95-102.

Das S., Seema P., Kundra C.P. and Pereira M.J., (2004): Reproduction in male rats is vulnerable to treatment with the flavanoid -rich seed extract of *Vitex negundo*. Phytotherapy Research 18, 8-13.

Gupta R.S., Yadav R., Dixit V.P. and Dobhal M.P., (2001): Antifertility studies of *Colebro oppositifolia* leaf extract in male rats special reference to cell population dynamics. Fitoterapia 73, 236-245.

Gupta R.S., Sharma R., Sharma A., Bhatnagar A., Dhobal M.P., Joshi Y.C. and Sharma M. G. (2002): Effect of *Alstonia scloris* bark extract on testicular function of wistar rats. Asian journal .of Andrology, 4 (3) 175-8.

Indian National Science academy (2000): Guidelines for care and use of animals in scientific research reported by Indian national science academy, New Delhi.

Ipstein J., Poly F. (1970): Brancroft's Introduction to biostatistics, IIInd Ed. (Harper International, New York) Pp. 44

Lowry O. H., Rosenberg N. J., Farr A. L. and Randall R. J., (1951): Protein measurement with the folin -phenol reagent. Journal of Biological Chemistry 193, 265-275

Lynch J. M., Raphel S. S., Melvir L. D., Spare P.D. and Inwood M. J. .M (1969): In medical laboratory and clinical pathology pub., Sounders Company SohmlTD., Tokyo 626, 647-662 .

Makrem A. (1974): Haemoglobin, myoglobin and hepatoglobin. In (Henry, Cannon,

- Winkelmann. Ed) Clinical Chemistry. Principles and techniques Pp. 1111-1214 (Harper and Row, London)
- WHO: Protocol cg-04. Preparation of alcoholic extract for bioassay and phytochemical studies (APJF/IP, 1001A) Geneva, World Health Organization 1983a.
- Montgomery R., (1957) Determination of glycogen. Arch Biochemistry and Biophysics 67, 378.
- Oser B. L., (1965): Hawk's physiological Chemistry. 14th Ed., New York, McGraw Hill, p 246.
- Sarkar M., Gangopadhyay P., Basak B., Chakrabarti K., Banerjee J., Adhikari P. and Chatterjee A., (2000): The reversible antifertility effect of *Piper beetle* Linn. On swiss albino mice *Contraception* 62, 271-274
- Sharma N. and Jacob D., (2001): Antifertility investigation and toxicological screening of the petroleum ether extract of the leaves of *Mentha arvensis* in male albino mice. Journal of Ethnopharmacology 75, 5-12.
- Shrikanth V., Malini T., Arunakaran J., Govindrajulu P., Balasubramanian K., (1999): Effect of ethanol treatment on epididymal secretary product and sperm maturation in albino rats. Journal of Pharmacology and Experimental Therapeutics 288, 509-515.
- Schalm O.W., Jain N.C., Carrolt E.J (1975): Veterinary haematology 3rd edition Lea and febiger Philadelphia PP 324_335
- Upadhyay S. Shanbhag K.K., Sunita G., and Balachandra Naidu., (2004): A study of hypoglycemic and antioxidant activity of *Aegle marmelos* in allonan induced diabetic rats . Indian Journal of physiological phamocology 48, 476-80
- Varley H (1969): Determination of blood urea by urease nesslerization method. In practical clinical biochemistry 4th edition. White Herres press .Ltd., London P 158-160
- Warren (1959): The thiobarbituric acid assay acid. Journal of Biological Chemistry 234, 1971.
- World Health Organization (1990): Special programme of research development and research training in human reproduction. Biennial report (1988-89) World health organization, Geneva.
- Srikanth V, Malini T, Arunakaran J, Govindarajulu P, Balasubramanian K. Effects of ethanol treatment on epididymal secretary products and sperm maturation in albino rats. J Pharmacol Exp Ther 1999; 288: 509-515.
- Zaneveld LJD, Polakoski KL. Collection and physical examination of the ejaculate. In: Hafez ESE, Ed., Techniques of Human Andrology. Amsterdam, Holland: north Biomedical Press, 1977. p. 147-156.
- Krag K. Plant used as contraceptive by the North American Indian :an ethanobotanical study. Botanical museum. Cambridge, MA: Harvared university, 1976: 1177.

Table 1: Effect of aqueous extract of *Calendula officinalis* (COFTAq) on body and reproductive organ weights of adult male rat

Treatment	Body Weight (gm)		Reproductive organs weight (mg/100gm)					vital organs weight (mg/100gm)		
	Initial	Final	Testis	Epididymis	Seminal Vesicle	Vas difference	Ventral prostate	Heart	kidney	liver
Group- I Control	185.30±08.01	204.20±07.20	1021.40±60.20	426.60±21.56	519.60±26.56	151.960±1.51	294.29±01.20	450.98±8.96	70592±1.20	4.9561±180.21
Group- II <i>Calendula officinalis</i> (100 -mg/kg b. wt.) per day for 60 days	180.28±2.03	200.00±5.26	939.67*±62.21	412.5*±26.69	500.560±26.69*	130.560±2.61 *	254.66±02.10*	440.96±7.98*	696.52±1.69*	4.894±176.01*

(Mean±SEM of 10 animals)

Group II compared with Group

Ins = Non-significant

** = P<0.001 - Highly Significant

Table 2: Effect of *Calendula officinalis* (COFTAq) on the sperm dynamics, fertility test and serum testosterone level male rat

Treatment	No. of males	No. of females	No. of pregnant females	No. of implantation sites	No. of viable fetuses	Sperm motility (Cauda epididymis) %	Sperm density (million/ml)		Fertility Test %	Serum Testosterone ng/dl
							Testes	Cauda epididymis		
Group- I Control	10	20	20/20	10.6±2.6	9.6±1.6	85.328±1.25	4.2±0.16	48.20±3.4	100(+)	4.71±0.12
Group- II <i>Calendula officinalis</i> (200- mg/kg b. wt.) per day for 60 days	10	20	11 /20	6.2± 2.3	8.63±1.27	78.526±1.26	3.02 ±0.43	43.10**±0.50	55(-)	1.87**±0.09

(Mean±SEM of 10 animals)

Group II compared with Group

Ins = Non-significant

* = P<0.01 - Significant

** = P<0.001 - Highly Significant

Table 3: Effect of *Calendula officinalis* (COFTAq) on the Tissue Biochemistry of adult male rat.

Treatment	Protein mg/100gm				Sialic Acid mg/100gm				Glycogen mg/100gm		Cholesterol mg/100gm	
	Testes	Epididymis	Seminal Vesicle	Ventral Prostate	Testes	Epididymis	Seminal Vesicle	Ventral Prostate	Testes	Liver	Testes	Liver
Group- I Control	245.16± 6.27	210.18± 5.17	195.18± 5.27	181.69± 0.26	5.89± 0.19	4.60± 0.67	5.86± 0.27	5.96± 0.69	3.78± 0.39	7.69± 1.43	7.99± 0.49	11.90± 2.31
Group- II <i>Calendula officinalis</i> (1000- mg/kg b. wt.) per day for 60 days	215.27± 6.19*	201.17±* 7.98	191.17*± 6.01	179.68± 1.28*	5.38± 0.26*	4.32.0± 0.89*	5.31± 0.67*	5.52 ±1.67 *	3.01 **± 0.52	6.65 **± 2.78	13.66± 2.92ns	8.46± 0.76ns

(Mean±SEM of 10 animals)

Group II compared

with Group I

ns = Non-significant

* = P<0.01 - Significant

** = P<0.001 - Highly Significant

Table 4: Effect of aqueous extract of *Calendula officinalis* (COFTAq) on haematological parameters of adult male rat

Treatment	Total Erythrocyte Count (TEC) million/mm ³	Total Leucocyte Count (TLC) per cu. Mm.	Total Leucocyte Count (TLC) per cu. Mm.	Haemoglobin gm%	Haematocrit % (pcv)	Blood sugar mg/100ml mg/dl	Blood Urea mg/dl mg/100ml
Group- I Control	5.02 ±0.62	8,602.3±406.9	8,603±406.9	14.0 ±0.39	45.23±2.6	96.86±0.23	96.86±0.23
Group- II <i>Calendula officinalis</i> (200-mg/kg b. wt.) per day for 60 days	5.06±0.69 ^{ns}	8,216.00 ^{ns} ±460.01	8,865.02 ^{ns} ±512.3	14.4 ^{ns} ±0.37 ^{ns}	45.26 ^{ns} ±6.8	95.87±0.69ns	39.02±3.8 ^{ns}

(Mean±SEM of 10 animals)

Group II compared with Group

Ins = Non-significant